



Europäisches Patentamt
European Patent Office
Office européen des brevets



Publication number: **0 520 372 A1**

EUROPEAN PATENT APPLICATION

Application number: **92110563.1**

Int. Cl.⁵: **C07F 9/40, A61K 31/66,
C07F 9/117**

Date of filing: **23.06.92**

Priority: **26.06.91 EP 91110546**

Date of publication of application:
30.12.92 Bulletin 92/53

Designated Contracting States:
**AT BE CH DE DK ES FR GB GR IT LI LU MC
NL PT SE**

Applicant: **HOECHST AKTIENGESELLSCHAFT
Postfach 80 03 20
W-6230 Frankfurt am Main 80(DE)**

Inventor: **Schiebler, Werner Dr.
Dresdener Strasse 5
W-6093 Flörsheim(DE)**

Inventor: **Deckert, Elke
Heimchenweg 86**

W-6230 Frankfurt am Main 80(DE)

Inventor: **Dreef, Eric Dr.c/o Gorlaeus
Laboratoria**

**Einsteinweg 5, Postbus 9502
2300 RA Leiden(NL)**

Inventor: **v.d.Marel, Geijs Dr., c/o Golaeus
Laboratoria**


**Einsteinweg 5, Postbus 9502
2300 RA Leiden(NL)**

Inventor: **von Boom, Jacques Prof., c/o
Gorlaeus Laboratoria**

**Einsteinweg 5, Postbus 9502
2300 RA Leiden(NL)**

Inositol phosphate analogues as calcium-antagonistic substances.

A compound of the formula (I)




in which

R₁ is e.g. phosphate,

R₂ is e.g. phosphate or OH, and

R₄ is e.g.

EP 0 520 372 A1



are suitable Calcium antagonists and useful e.g. for the treatment of cardio-vascular diseases.


The subject of this invention is a novel group of inositol-phosphat derivatives and their use as calcium-antagonistic compounds for different pharmacological uses such as for the treatment of cardio-vascular diseases.

It is now generally accepted that hydrolysis of phosphatidylinositol(4,5)bispophosphate (PIP2) by receptor- and G-protein mediated activation of PIP2-specific phospholipase C (PIP2-PLC) results in the formation of two different second messengers in the cytoplasm of cells: d-myo inositol(1,4,5)trisphosphate (IP3) and diacylglycerol. IP3 can bind to an appropriate receptor on intracellular organelles causing release of Calcium. Two metabolic pathways terminate this receptor-mediated Calcium release: a) the action of an IP3-3-kinase, phosphorylating the 3-position leading to Inositol (1,3,4,5) tetrakisphosphate (IP4) and b) the action of an IP3-5-phosphatase leading to Inositol(1,4)bispophosphate. Since the cell is expending energy in form of ATP to generate IP4, IP4 is considered to be a putative second messenger, possibly regulating further steps in the Calcium-Metabolism of cells (M.J. Berridge, (1987) Ann. Rev. Biochem. 56, 159-193; and M.J. Berridge and R.F. Irvine, (1989) Nature 341, 197-205). The importance of IP3 as a Calcium mobilizing intracellular second messenger has lead to the synthesis of IP3 and derivatives thereof.

Preparation of Phosphorothioates acting as long-lived agonists have been described (A.M. Cooke et al., (1987) J. Chem. Soc. Chem. Commun. 1525-1526; and J. Strupish et al., (1988) Biochem. J. 253, 901-905). However, there have been no reports in IP3-analogues acting as IP3-receptor- and therefore intracellular Calcium-antagonists. In addition, analogues of IP4 acting as long lived agonists or antagonists, are not available yet.

It is an object of this invention, to describe the synthesis of specific IP3-, IP4- and PIP/PIP2-Analogues modified at the 5-Position. A further surprising aspect of this invention is to describe antagonists of the IP3-receptor, acting as intracellular Calcium-antagonists. These substances can generally be used as highly active pharmacological compounds, especially they can be used as antithrombotic agents and in a variety of other cardiovascular diseases.

Subject of this invention is especially a compound of the formula (I)



in which
R1 is

- 1) phosphate
- 2) OH
- 3) 1,2 diacyl-sn-glycer-3-yl phosphate, wherein acyl is a fatty acid with 2 to 20 C-atoms and 0 to 4 C-C-double bonds, or
- 4) 1,2 dialkyl-sn-glycer-3-yl phosphate, wherein alkyl is a hydrocarbon having 2 to 20 carbon atoms and 0 to 4 C-C-double bonds

R2 is

- 1) OH, or
- 2) phosphate

R3 is

- 1) phosphate, or
- 2) OH
- 3) a radical of formula (II)



wherein

- a) Y = Z = O and X = (C₁-C₈)-alkyl
 - b) Y = Z = O and X = (C₆-C₁₈)-aryl
 - c) Y = S and Z = O and X = (C₁-C₈)-alkyl
 - d) Y = S and Z = O and X = (C₆-C₁₈)-aryl
 - e) Y = O and Z = X = (C₁-C₈)-alkyl
 - f) Y = O and Z = X = (C₆-C₁₈)-aryl
 - g) Y = S and Z = X = (C₁-C₈)-alkyl
 - h) Y = S and Z = X = (C₆-C₁₈)-aryl
 - i) Y = Z = O and X = CF₂H
 - j) Y = O and Z = X = F
 - k) Y = Z = O and X = O-(C₁-C₈)-alkyl
 - l) Y = Z = O and X = O-(C₆-C₁₈)-aryl
 - m) Y = S and Z = O and X = O-(C₁-C₈)-alkyl
 - n) Y = S and Z = O and X = O-(C₆-C₁₈)-aryl
 - o) Y = O and Z = X = O-(C₁-C₈)-alkyl
 - p) Y = O and Z = X = O-(C₆-C₁₈)-aryl
 - q) Y = S and Z = X = O-(C₁-C₈)-alkyl
 - r) Y = S and Z = X = O-(C₆-C₁₈)-aryl
- 4) a radical of formula (III)



where

- a) X = OR where R is H, (C₁-C₈)-alkyl, or (C₆-C₁₈)-aryl
 - b) X = (C₁-C₈)-alkyl
 - c) X = (C₆-C₁₈)-aryl
 - d) X = NH-(C₁-C₈)-alkyl
 - e) X = NH-(C₆-C₁₈)-aryl
- 5) a radical of formula (IV)




where

- a) X = OR where R is H, (C₁-C₈)-alkyl or (C₆-C₁₈)-aryl
- b) X = (C₁-C₈)-alkyl
- c) X = (C₆-C₁₈)-aryl
- d) X = NH-(C₁-C₈)-alkyl
- e) X = NH-(C₆-C₁₈)-aryl

R₄ is

- 1) a radical of formula (V)



where

- a) Y = Z = O and X = (C₁-C₈)-alkyl
 - b) Y = Z = O and X = (C₆-C₁₈)-aryl
 - c) Y = S and Z = O and X = (C₁-C₈)-alkyl
 - d) Y = S and Z = O and X = (C₆-C₁₈)-aryl
 - e) Y = O and Z = X = (C₁-C₈)-alkyl
 - f) Y = O and Z = X = (C₆-C₁₈)-aryl
 - g) Y = S and Z = X = (C₁-C₈)-alkyl
 - h) Y = S and Z = X = (C₆-C₁₈)-aryl
 - i) Y = Z = O and X = CF₂H
 - j) Y = O and Z = X = F
 - k) Y = Z = O and X = O-(C₁-C₈)-alkyl
 - l) Y = Z = O and X = O-(C₆-C₁₈)-aryl
 - m) Y = S and Z = O and X = O-(C₁-C₈)-alkyl
 - n) Y = S and Z = O and X = O-(C₆-C₁₈)-aryl
 - o) Y = O and Z = X = O-(C₁-C₈)-alkyl
 - p) Y = O and Z = X = O-(C₆-C₁₈)-aryl
 - q) Y = S and Z = X = O-(C₁-C₈)-alkyl
 - r) Y = S and Z = X = O-(C₆-C₁₈)-aryl
- 2) a radical of formula (VI)



wherein

- a) X = OR where R is H, (C₁-C₈)-alkyl, or (C₆-C₁₈)-aryl
 - b) X = (C₁-C₈)-alkyl
 - c) X = (C₆-C₁₈)-aryl
 - d) X = NH-(C₁-C₈)-alkyl
 - e) X = NH-(C₆-C₁₈)-aryl
- 3) a radical of formula (VII)



wherein

- a) X = OR where R is H, (C₁-C₈)-alkyl or (C₆-C₁₈)-aryl
- b) X = (C₁-C₈)-alkyl
- c) x = (C₆-C₁₈)-aryl
- d) X = NH-(C₁-C₈)-alkyl
- e) X = NH-(C₆-C₁₈)-aryl


and the physiologically tolerated salts thereof.

Preferred are those compounds of the formula (I) in which

R1 is phosphate or 1,2-diacyl-sn-glycer-3-yl-phosphate, wherein acyl is a fatty acid radical having 2 to 20 carbon atoms and 0 to 4 C-C-double bonds,

R2 is OH or phosphate,

- R3 is OH or phosphate,
 R4 is a radical of formula (V),



10 wherein Y = Z = O and X = (C₁-C₈)-alkyl, and the physiologically tolerated salts thereof.

Furthermore the radical R⁴ has the following preferred definition:

R⁴ is a radical of the formula (V) wherein X is methyl, ethyl or propyl.

Another part of the invention is the general synthesis of inositol phosphate derivatives of the formula (I) starting with myo-inositol derivatives like for instance a 3,6-di-O-allyl-1-2-O-cyclohexylidene-myo-inositol (1),
 15 proceeding through synthesis of e.g. 3,6-di-O-allyl-4-O-benzyl-1,2-O-cyclohexylidene-myo-inositol (2b) (the respective numbering using the 1-position of the natural Inositol(1.4.5)trisphosphate would be 1,4-di-O-allyl-6-O-benzyl-2,3-O-cyclohexylidene-myo-inositol). Through a series of protection and deprotection steps which are generally known to a person skilled in this field, using protection of the 5-position with e.g. a -O-p-methoxybenzyl-group, 5-position analogues of both Inositol(1.4.5)trisphosphate and Inositol(1.3.4.5)-
 20 tetrakisphosphate can be synthesized.

The compounds of formula (I) can act as calcium-antagonists.

Accordingly, these compounds (inositol phosphate analogues) administered in the appropriate formulation and appropriate dosis, can act as different pharmacological agents, e.g. as antithrombotic agents. Since contraction of smooth muscles also depend on intracellular release of Calcium, vasorelaxant effects of the
 25 compounds can be used to treat diseases like hypertension. General calcium-antagonistic effects of the compounds can be exploited to treat diseases like Asthma, Cardiomyopathy, Atherosclerosis, Ischemia (Heart and Brain) and Inflammatory Diseases.

The compounds can be applied in the appropriate dosis, preferably 0.01-100 mg/kg, either orally, intranasally, intraperitoneally, intravenously, topically, or subcutaneously. Dosis and application depends on
 30 the species, preferably mammals, bodyweight and age.

Pharmaceutical preparations contain an effective amount of the compound together with a suitable vehicle like e.g. a compatible inorganic or organic carrier. The usual additions for the respective oral, intravenous, topical etc. applications can be used.

The invention is further illustrated by the following examples. In the experimental part the synthesis is
 35 outlined by the following schemes A, B and C.


40

45


50

55


Scheme A



Scheme B




Scheme C



Example 1

50 Synthesis of myo-inositol 5-methylphosphonate 1,4-bisphosphate



a) 3,6-di-O-allyl-4,5-di-O-benzyl-1,2-O-cyclohexylidene-myo-inositol (2a), 3,6-di-O-allyl-4-O-benzyl-1,2-O-cyclohexylidene-myo-inositol (2b) and 3,6-di-O-allyl-5-O-benzyl-1,2-O-cyclohexylidene-myo-inositol (2c)

To a solution of compound 1, (see C.E. Dreef et al., Recl. Trav. Chim Pays-Bas, 106, 161 (1987)), (6.80 g, 20.00 mmol) and tetrabutylammonium hydrogen sulfate (6.78 g, 20.00 mmol) in CH_2Cl_2 (200 mL) were added benzyl bromide (3.00 mL, 25.23 mmol) and 5 % aqueous NaOH (200 mL). The reaction mixture was refluxed for 24 h. The organic layer was separated and washed with H_2O , 1 M NaHCO_3 and H_2O . The organic layer was dried over MgSO_4 and concentrated in vacuo. Silica gel column chromatography (140 g, elution: hexane/ Et_2O , 100/0 to 50/50, v/v) of the crude product mixture afforded :

Compound 2a (0.52 g, 5 % yield) as an oil;

$^1\text{H-NMR}$ (CDCl_3): delta 1.27-1.80 (m, 10H, 5 x CH_2 , cyclohexylidene), 3.37 (dd, 1H, H-5, $J_{5,6} = 9.5$ Hz), 3.63 (dd, 1H, H-3, $J_{3,4} = 8.5$ Hz), 3.68 (dd, 1H, H-6, $J_{6,1} = 7.0$ Hz), 3.87 (dd, 1H, H-4, $J_{4,5} = 8.5$ Hz), 4.07 (dd, 1H, H-1, $J_{1,2} = 5.5$ Hz), 4.18-4.29 (m, 3H, 2 x OCH_2 , All), 4.34-4.43 (m, 1H, 2 x OCH_2 , All), 4.39 (dd, 1H, H-2, $J_{2,3} = 4.0$ Hz), 4.77-4.85 (m, 4H, 2 x OCH_2 , Bzl), 5.14-5.20 (m, 4H, 2 x $=\text{CH}_2$, All), 5.89-6.05 (m, 2H, 2 x $-\text{CH} =$, All), 7.25-7.38 (m, 10H, CH, H aromatic).

Compound 2b (4.56 g, 53 % yield) as an oil;

$^1\text{H-NMR}$ (CDCl_3): delta 1.25-1.82 (m, 10H, 5 x CH_2 , cyclohexylidene), 2.69 (d, 1H, 5-OH (exchangeable)) 3.46 (ddd, 1H, H-5, $J_{5,6} = 9.5$ Hz, $J_{5,\text{OH}} = 2.0$ Hz), 3.60 (dd, 1H, H-6, $J_{6,1} = 7.0$ Hz), 3.64 (dd, 1H, H-3, $J_{3,4} = 8.0$ Hz), 3.76 (dd, 1H, H-4, $J_{4,5} = 8.0$ Hz), 4.07 (dd, 1H, H-1, $J_{1,2} = 5.5$ Hz), 4.19-4.25 (m, 3H, 2 x OCH_2 , All), 4.37-4.43 (m, 1H, 2 x OCH_2 , All), 4.41 (dd, 1H, H-2, $J_{2,3} = 4.0$ Hz), 4.77-4.91 (m, 2H, OCH_2 , Bzl), 5.17-5.33 (m, 4H, 2 x $=\text{CH}_2$, All), 5.90-6.01 (m, 2H, 2 x $-\text{CH} =$, All), 7.25-7.41 (m, 5H; H aromatic), and compound 2c (3.10 g, 36 % yield) as an oil;

$^1\text{H-NMR}$ (CDCl_3): delta 1.26-1.76 (m, 10H, 5 x CH_2 , cyclohexylidene), 2.63, (d, 1H, 4-OH (exchangeable)) 3.25 (dd, 1H, H-5, $J_{5,6} = 9.0$ Hz), 3.50 (dd, 1H, H-3, $J_{3,4} = 10.0$ Hz), 3.60 (dd, 1H, H-6, $J_{6,1} = 7.0$ Hz), 3.97 (ddd, 1H, H-4, $J_{4,5} = 9.0$ Hz, $J_{4,\text{OH}} = 1.5$ Hz), 4.08 (dd, 1H, H-1, $J_{1,2} = 5.0$ Hz), 4.19-4.26 (m, 3H, 2 x OCH_2 , All), 4.33-4.40 (m, 1H, 2H, OCH_2 , Bzl), 5.15-5.35 (m, 4H, 2 x $=\text{CH}_2$, All), 5.89-6.02 (m, 2H, 2 x $-\text{CH} =$, All), 7.25-7.42 (m, 5H, H aromatic).

b) 1,4-di-O-allyl-6-O-benzyl-5-O-p-methoxybenzyl-myo-inositol (4)

To a solution of compound 2b (4.30 g, 10.00 mmol) and NaH (0.30 g, 12.50 mmol) in dry DMF (50 ml) was added dropwise p-methoxybenzyl chloride (1.50 ml, 11.07 mmol) at 0 °C. The reaction mixture was stirred for 2 h at 20 °C. Excess NaH was destroyed with MeOH and the reaction mixture concentrated in vacuo. The residue was taken up in CH_2Cl_2 , washed with H_2O , 1 M NaHCO_3 and H_2O . The organic layer was dried over MgSO_4 and concentrated in vacuo. To a solution of the crude compound 3 in MeOH (25 ml) was added 0.1 N HCl in MeOH (25 ml, 2.50 mmol) and the reaction mixture was stirred for 5 h at 20 °C. The reaction mixture was neutralized with Et_3N and concentrated in vacuo. The residue was taken up in CH_2Cl_2 , washed with H_2O , 1 M NaHCO_3 and H_2O . The organic layer was dried over MgSO_4 and concentrated in vacuo. Silica gel column chromatography (60 g, solution: $\text{CH}_2\text{Cl}_2/\text{MeOH}$, 100/0 to 95/5, v/v) of the crude product afforded pure 4 (4.09 g, 87 % yield), Mp 97-98 °C (from $\text{Et}_2\text{O}/\text{hexane}$).

$^1\text{H-NMR}$ (CDCl_3): delta 2.52 (s (br), 1H, 2-OH, exchangeable), 2.55 (d, 1H, 3-OH, exchangeable), 3.36 (dd, 1H, H-1, $J_{1,2} = 3.0$ Hz), 3.39 (dd, 1H, H-5, $J_{5,6} = 9.5$ Hz), 3.46 (ddd, 1H, H-3, $J_{3,4} = 9.5$ Hz, $J_{3,\text{OH}} = 4.5$ Hz), 3.68 (dd, 1H, H-4, $J_{4,5} = 9.5$ Hz), 3.80 (s, 3H, OCH_3 , pMeOBzl), 3.86 (dd, 1H, H-6, $J_{6,1} = 9.5$ Hz), 4.13-4.29 (m, 3H, 2 x OCH_2 , All), 4.22 (dd (br), 1H, H-2, $J_{2,3} = 3.0$ Hz, $J_{2,\text{OH}} < 1.0$ Hz), 4.40-4.47 (m, 1H, 2 x OCH_2 , All), 4.70-4.89 (m, 4H, 2 x OCH_2 , Bzl and pMeOBzl), 5.17-5.34 (m, 4H, 2 x $=\text{CH}_2$, All) 5.87-6.03 (m, 2H, 2 x $-\text{CH} =$, All), 6.82-6.87 and 7.21-7.37 (m, 9H, H aromatic). A separation of the enantiomers may be achieved e.g. by chiral auxiliary groups.

c) 1,4-di-O-allyl-2,3,6-tri-O-benzyl-5-O-p-methoxybenzyl-myo-inositol (5)

To a solution of compound 4 (1.75 g, 3.72 mmol) and NaH (0.23 g, 9.58 mmol) in dry DMF (20 mL) was added dropwise benzyl bromide (1.00 ml, 8.41 mmol) at 0 °C. The reaction mixture was stirred for 2 h at 20 °C. Excess NaH was destroyed with MeOH and the reaction mixture concentrated in vacuo. The residue was taken up in CH_2Cl_2 , washed with H_2O , 1 M NaHCO_3 and H_2O . The organic layer was dried over MgSO_4 and concentrated in vacuo. Silica gel column chromatography (30 g, elution: hexane/ Et_2O , 100/0 to 50/50, v/v) of the crude product yielded 5 (2.30 g, 95 % yield, Mp 69.5-70.5 °C (from pentane)).

$^1\text{H-NMR}$ (CDCl_3): delta 3.23 (dd, 1H, H-1, $J_{1,2} = 2.5$ Hz), 3.30 (dd, 1H, H-3, $J_{3,4} = 10.0$ Hz), 3.38 (dd, 1H, H-5, $J_{5,6} = 9.0$ Hz), 3.79 (s, 3H, OCH_3 , pMeOBzl), 3.91 (dd, 1H, H-4, $J_{4,5} = 9.0$ Hz), 3.96 (dd, 1H, H-6, $J_{6,1}$

= 10.0 Hz), 3.99 (dd, 1H, H-2, $J_{2,3}$ 2.5 Hz), 4.03-4.13 (m, 2H, OCH₂, All) 4.29-4.44 (m, 2H, OCH₂, All), 4.58-4.90 (m, 8H, 4 x OCH₂, Bzl and pMeOBzl), 5.13-5.32 (m, 4H, 2 x =CH₂, All), 5.83-6.06 (m, 2H, 2 x -CH=, All), 6.82-6.86 and 7.22-7.41 (m, 19H, H aromatic).

5 d) 2,3,6-tri-O-benzyl-5-O-p-methoxybenzyl-myo-inositol (6)

To a solution of compound 5 (1.95 g, 3.00 mmol) in 1,2-dichloroethane (15 ml) under an inert helium atmosphere was added 1,5-cyclooctadiene-bis[methyldiphenylphosphine]iridium hexafluorophosphate (see L.M. Haines et al., J. Chem. Soc. Dalton Trans., 1891 (1972)), (20 mg) in 1,2-dichloroethane (0.5 ml). The catalyst was activated by passing over a stream of hydrogen for 2 min. The solution was degassed and left under a stream of argon for 4 h. The reaction mixture was concentrated in vacuo and the crude 2,3,6-tri-O-benzyl-5-O-p-methoxybenzyl-1,4-di-O-trans-prop-1-enyl-myo-inositol, thus obtained was used without further purification.

To a solution of the crude 2,3,6-tri-O-benzyl-5-O-p-methoxybenzyl-1,4-di-O-trans-prop-1-enyl-myo-inositol in CH₂Cl₂ (15 ml) was added 0.2 N HCl in MeOH (15 ml, 3.00 mmol) and the reaction mixture was stirred for 1 h at 20 °C. The reaction mixture was neutralized with Et₃N and concentrated in vacuo. The residue was taken up in CH₂Cl₂, washed with H₂O, 1 M NaHCO₃ and H₂O. The organic layer was dried over MgSO₄ and concentrated in vacuo. Silica gel column chromatography (25 g, elution: CH₂Cl₂/acetone, 100/0 to 97/3, v/v) of the crude product afforded pure 6 (1.58 g, 92 % yield). Mp 97.5-98.5 °C (from Et₂O/hexane).

¹H-NMR (CDCl₃): delta 2.28 (d, 1H, 1-OH, exchangeable), 2.52 (d, 1H, 4-OH, exchangeable), 3.30 (dd, 1H, H-3, $J_{3,4}$ = 10.0 Hz), 3.38 (dd, 1H, H-5, $J_{5,6}$ = 9.0 Hz), 3.52 (ddd, 1H, H-1, $J_{1,2}$ = 2.5 Hz; $J_{1,OH}$ = 6.5 Hz), 3.78 (dd, 1H, H-6, $J_{6,1}$ = 9.5 Hz), 3.80 (s, 3H, OCH₃, pMeOBzl), 4.07 (dd, 1H, H-2, $J_{2,3}$ = 2.5 Hz), 4.14 (ddd, 1H, H-4, $J_{4,5}$ = 9.0 Hz, $J_{4,OH}$ = 1.5 Hz), 4.59-4.94 (m, 8H, 4 x OCH₂, Bzl and pMeOBzl), 6.84-6.89 and 7.21-7.40 (m, 19H, H aromatic).

e) 2,3,6-tri-O-benzyl-5-O-p-methoxybenzyl-myo-inositol 1,4-bis-(dibenzylphosphate) (10)

A mixture of compound 6 (1.43 g, 2.51 mmol) and N,N-diisopropylidibenzyl phosphoramidite (see K.-L. Yu and B. Fraser-Reid, Tetrahedron Let., 29, 979 (1988)) (2.60 g, 7.54 mmol) was coevaporated with toluene (2 x 25 ml) and dissolved in CH₂Cl₂ (20 ml). Subsequently a solution of 1H-tetrazole (0.65 g, 9.29 mmol) in CH₃CN (20 ml) was added and the reaction mixture was stirred for 15 min. ³¹P-NMR showed the presence of two peaks (delta 141.04 and 141.88). The reaction mixture was cooled (0 °C) and tert-butyl hydroperoxide (3.75 ml) was added and stirring was continued for 45 min at 0 °C. The reaction mixture was diluted with CH₂Cl₂ and washed with H₂O. The organic layer was dried over MgSO₄ and concentrated in vacuo. Silica gel column chromatography (35 g, elution: hexane/EtOAc, 100/0 to 25/75, v/v) of the crude product afforded homogeneous 10 (2.57 g, 94 % yield) as an oil.

³¹P-NMR (CH₂Cl₂): delta -1.03 (2P)

40 f) 2,3,6-tri-O-benzyl-myo-inositol 1,4-bis(dibenzylphosphate) (11)

To a solution of compound 10 (2.18 g, 2.00 mmol) in CH₂Cl₂ (48.75 ml) was added trifluoroacetic acid (1.25 ml) and the reaction mixture was stirred for 30 min at 20 °C. The reaction mixture was diluted with CH₂Cl₂ and washed with H₂O, 1 M Tetraethylammonium bicarbonate buffer (TEAB) and H₂O. The organic layer was dried over MgSO₄ and concentrated in vacuo. Silica gel column chromatography (25 g, elution: hexane/EtOAc, 100/0 to 25/75, v/v) of the crude product afforded pure 11 (1.59 g, 82 % yield), as an oil.

³¹P-NMR (CH₂Cl₂): delta -1.24(1P) and 0.76 (1P).

g) 2,3,6-tri-O-benzyl-myo-inositol 5-(benzyl methylphosphonate) 1,4-bis(dibenzylphosphate) (13)

A solution of bis(1-[6-trifluoromethyl]benzotriazolyl) methylphosphonate in dioxane (0.2 M, 3.5 ml, 0.70 mmol) was added to compound 11 (0.34 g, 0.35 mmol), which had been dried by repeated coevaporation with pyridine. The reaction was stirred for 30 min at 20 °C. Subsequently benzyl alcohol (0.15 ml, 1.45 mmol) and N-methylimidazole (0.15 ml, 1.88 mmol) were added and the reaction mixture was stirred for another 1 h at 20 °C. After addition of 1 M TEAB the reaction mixture was diluted with CH₂Cl₂ and washed with H₂O, 1 M TEAB and H₂O. The organic layer was dried over MgSO₄ and concentrated in vacuo. Silica gel column chromatography (5 g, elution: hexane/EtOAc, 100/0 to 25/75, v/v) of the crude product afforded pure 13 (0.30 g, 75 % yield), as an oil.

^{31}P -NMR (CH_2Cl_2): δ -1.27(2P); 31.31 and 32.88 (1P, ratio, 1/3).

h) myo-inositol 5-methylphosphonate 1,4-bisphosphate (Na^+ -form) (14)

Compound 13 (225 mg, 0.20 mmol) was dissolved in a mixture of MeOH and H_2O (50 ml, 4/1, v/v) and hydrogenated over 10 % palladium on charcoal (0.20 g) at 500 kPa for 16 h at 20 °C. The solution was filtered and concentrated in vacuo (30 °C) to a small volume. After Sephadex C-25 (Na^+ -form, 4.5 g, 10.4 mmol) cation-exchange and lyophilization 14 (101 mg, 97 % yield) was obtained, as a white solid.

^{31}P -NMR (D_2O , pH = 2.00): δ 0.29 (P-1), 1.20 (P-4) and 31.68 (P-5).

^1H -NMR (D_2O , pH = 2.00): δ 1.48 (d, 3H, CH_3 , $J_{\text{H,P}} = 17.5$ Hz), 3.75 (dd, 1H, H-3, $J_{3,4} = 10.0$ Hz), 3.88 (dd, 1H, H-6, $J_{6,1} = 10.0$ Hz), 4.03 (ddd, 1H, H-1, $J_{1,2} = 2.5$ Hz, $J_{\text{H,P}} = 8.5$ Hz), 4.12 (ddd, 1H, H-5, $J_{5,6} = 9.5$ Hz, $J_{\text{H,P}} = 9.0$ Hz), 4.28 (dd, 1H, H-2, $J_{2,3} = 3.0$ Hz), 4.34 (ddd, 1H, H-4, $J_{4,5} = 9.0$ Hz, $J_{\text{H,P}} = 9.0$ Hz).

Example 1b

Pharmacological activity of the compound of example 1

The activity of myo-inositol 5-methylphosphonate 1,4-bisphosphate (IP2-5-MP) as a Calcium antagonist in human platelets is shown as follows.

Human platelets were purified in the presence of citrate by differential centrifugation and resuspended to ca 1×10^9 cells/ml in Tyrode-HEPES-(N-2-hydroxyethylpiperazine-N'-2 ethane-sulfonic acid)-buffer pH 7.4 containing 0.2 % w/v Bovine Serum Albumin and 1mM EGTA (Ethylene Glycol Bis (β -Aminoethyl Ether)N,N,N',N'-Tetraacidic acid). After 20 min, the platelets were centrifuged again (700xg, 20 min) and resuspended to the same concentration in 4 ml of "assay-buffer" containing:


110 mM	Potassiumchloride
1.2 mM	Potassiumdihydrogenphosphate
5 mM	Potassiumsuccinate
5 mM	Potassiumpyruvate
5 mM	Potassium-ATP
6.8 mM	Magnesiumchloride
9.8 U/ml	Creatine kinase
12 mM	Creatine phosphate
25 mM	HEPES-buffer, pH as indicated

After a further 30 min, the platelet suspension was diluted 5-fold into assay-buffer containing 2 μM Quin2 (2-[[2-[bis(carboxymethyl)amino]-5-methyl-phenoxy]methyl]-6-methoxy-8-[bis(β -aminoethyl)amino]-quinoline) salt as a fluorescence indicator for Calcium. To this suspension Saponin was added to a final concentration of 20 $\mu\text{g}/\text{ml}$ to permeabilize the cells. The fluorescence of the suspension was continually monitored with a fluorescence spectrometer until a stable baseline was achieved. At this point, IP3 was added and the resulting release of Calcium measured as an increase in the intensity of the Quin2-fluorescence. An EC50 (effective concentration) value of 2×10^{-7} M was obtained at pH 7.4. IP2-5-MP itself had no effect on release of Calcium. However, when applied at 10^{-4} M 15 sec before addition of 5×10^{-7} M IP3, at pH values below 7.4 the Calcium response to IP3 was inhibited.

The following Inhibitor-values were found at the indicated concentrations of IP3 (5×10^{-7} M) and IP2-5-MP (1×10^{-4} M):


pH	Inhibition of Calcium release (%)
7.6	0
7.4	0
7.3	68
7.2	100
7.1	100

Example 2

Synthesis of myo-inositol 5-ethylphosphonate 1,4-bisphosphate (Na⁺-form)

15 To compound 11 a solution of bis(1-[6-trifluoromethyl]ethyl)phosphonate was added under identical conditions as for compound 13. Following the amounts and reaction conditions as given for the compounds 13-14, the final product myo-inositol 5-ethylphosphonate 1,4-bisphosphate (Na⁺-form) was synthesized. ³¹P-NMR-Signals: (D₂O): delta-3.54 (P-1 and P-4), 32,70 (P-5).

20 Example 3

Synthesis of myo-inositol 5-((difluoromethyl)phosphonate) 1,4-bis-phosphate (Na⁺-form)

a) 2,3,6-tri-O-benzyl-myo-inositol 5-(benzyl(difluoromethyl)phosphonate) 1,4-bis(dibenzylphosphate) (15)

35 A solution of (difluoromethyl)phosphonic di(1,2,4-triazolide) in dioxane (0.2 M, 3.5. ml, 0.780 mmol) was added to compound 11(0.34 g, 0.35 mmol), which had been dried by repeated coevaporation with pyridine. The reaction was stirred for 30 min at 20 °C. Subsequently benzyl alcohol (0.15 ml, 1.45 mmol) and N-methylimidazole (0.15 ml, 1.88 mmol) were added and the reaction mixture was stirred for another 1 h at 20 °C. After addition of 1 M TEAB the reaction mixture was diluted with CH₂Cl₂ and washed with H₂O, 1 M TEAB and H₂O. The organic layer was dried over MgSO₄ and concentrated in vacuo. Silica gel column chromatography (5 g, elution: Hexane/EtOAc, 100/0 to 25/75, v/v) of the crude product afforded pure 15 (0.29 g, 70 % yield), as an oil.

40 ³¹P-NMR (CH₂Cl₂: delta -1.45, -1.36, -1.18 and -1.00(2P); 4.32 (JP,F = 93.0 Hz) and 6.66 (JP,F = 85.5 Hz and JP,F = 97.5 Hz) (1P).


45 b) myo-inositol 5-((difluoromethyl)phosphonate) 1,4-bisphosphate (Na⁺-form) (16)

Compound 15 (220 mg, 0.19 mmol) was dissolved in a mixture of MeOH and H₂O (50 ml, 4/1, v/v) and hydrogenated over 10 % palladium on charcoal (0.20 g) at 500 kPa for 16 h at 20 °C. The solution was filtered and concentrated in vacuo (30 °C) to a small volume. After Sephadex C-25 (Na⁺-form, 4.0 g, 9.2 mmol) cation-exchange and lyophilization 16(0.10 g, 95% yield) was obtained, as a white solid.

50 ³¹P-NMR (D₂O, pH = 2.00): delta 0.27 (P-1), 1.16 (P-4) and 4.68 (P-5, JP,F = 85.0 Hz).

Example 4

55 Synthesis of myo-inositol 5-methylphosphonate 1,3,4-trisphosphate (Na⁺-form)



a) 1,3,4-tri-O-allyl-6-O-benzyl-5-O-p-methoxybenzyl-myoinositol (7)

A solution of compound 4 (2.00 g, 4.26 mmol) and dibutyltin oxide (1.20 g, 4.82 mmol) in dry MeOH (25 ml) was refluxed for 2.5 h and subsequently concentrated in vacuo. The residue was coevaporated with toluene (3 x 25 ml), dissolved in dry DMF (45 ml) and subsequently cesium fluoride (0.85 g, 5.59 mmol) and allyl bromide (0.55 mL, 6.50 mmol) were added. After stirring for 16 h at 20 °C the reaction mixture was concentrated in vacuo. The residue was taken up in Et₂O, washed with H₂O, 1 M NaHCO₃ and H₂O. The organic layer was dried over MgSO₄ and concentrated in vacuo. Silica gel column chromatography (30 g, elution: hexane/EtOAc, 100/0 to 50/50, v/v) of the crude product afforded pure 7 (1.81 g, 83 % yield). Mp 77-78 °C (from Et₂O/pentane).

¹H-NMR (CDCl₃): delta 2.42 (s (br), 1H, 2-OH, exchangeable), 3.25 (dd, 1H, H-3, J_{3,4} = 9.5 Hz), 3.29 (dd, 1H, H-1, J_{1,2} = 3.0 Hz), 3.36 (dd, 1H, H-5, J_{5,6} = 9.5 Hz), 3.77 (dd, 1H, H-4, J_{4,5} = 9.5 Hz), 3.80 (s, 3H, OCH₃, pMeOBzl), 3.87 (dd, 1H, H-6, J_{6,1} = 9.5 Hz), 4.18-4.22 (m, 4H, 2 x OCH₂, All), 4.21 (dd (br), 1H, H-2, J_{2,3} = 3.0 Hz), 4.25-4.39 (m, 2H, OCH₂, All), 4.72-4.87 (m, 4H, 2 x OCH₂, Bzl and pMeOBzl), 5.14-5.35 (m, 6H, 3 x =CH₂, All), 5.88-6.05 (m, 3H, 3 x -CH=, All), 6.82-6.87 and 7.23-7.38 (m, 9H, H aromatic).

b) 1,3,4-tri-O-allyl-2,6-di-O-benzyl-5-O-p-methoxybenzyl-myoinositol (8)

To a solution of compound 7 (1.75 g, 3.43 mmol) and NaH (0.11 g, 4.58 mmol) in dry DMF (20 ml) was added dropwise benzyl bromide (0.45 ml, 3.78 mmol) at 0 °C. The reaction mixture was stirred for 2 h at 20 °C. Excess NaH was destroyed with MeOH and the reaction mixture concentrated in vacuo. The residue was taken up in CH₂Cl₂, washed with H₂O, 1 M NaHCO₃ and H₂O. The organic layer was dried over MgSO₄ and concentrated in vacuo. Silica gel column chromatography (30 g, elution: hexane/Et₂O, 100/0 to 50/50, v/v) of the crude product yielded 8 (1.96 g, 95 % yield) as an oil. Mp 39.5-40.5 °C (solidified).

¹H-NMR (CDCl₃): delta 3.19 (dd, 1H, H-3, J_{3,4} = 10.0 Hz), 3.24 (dd, 1H, H-1, J_{1,2} = 2.5 Hz), 3.37 (dd, 1H, H-5, J_{5,6} = 9.0 Hz), 3.79 (s, 3H, OCH₃, pMeOBzl), 3.86 (dd, 1H, H-4, J_{4,5} = 9.0 Hz), 3.96 (dd, 1H, H-6, J_{6,1} = 10.0 Hz), 3.99 (dd, 1H, H-2, J_{2,3} = 2.5 Hz), 4.02-4.16 (m, 4H, 2 x OCH₂, All), 4.26-4.41 (m, 2H, OCH₂, All), 4.69-4.90 (m, 6H, 3 x OCH₂, Bzl and pMeOBzl), 5.13-5.34 (m, 6H, 3 x =CH₂, All), 5.85-6.06 (m, 3H, 3 x -CH=, All), 6.82-6.86 and 7.23-7.44 (m, 14H, H aromatic).

c) 2,6-di-O-benzyl-5-O-p-methoxybenzyl-myoinositol (9)

To a solution of compound 8 (1.80 g, 3.00 mmol) in 1,2-dichloro-ethane (15 ml) under an inert helium atmosphere was added 1,5-cyclooctadiene-bis[methyldiphenylphosphine]iridium hexafluoro-phosphate (see L.M. Haines et al., J. Chem. Soc. Dalton Tans. 1891 (1972)) (20 mg) in 1,2-dichloroethane (0.5 ml). The catalyst was activated by passing over a stream of hydrogen for 2 min. The solution was degassed and left under a stream of argon for 4 h. The reaction mixture was concentrated in vacuo and the crude 2,6-di-O-benzyl-5-O-p-methoxybenzyl-1,3,4-tri-O-trans-prop-1-enyl-myoinositol, thus obtained was used without further purification.

To a solution of the crude 2,6-di-O-benzyl-5-O-p-methoxybenzyl-1,3,4-tri-O-trans-prop-1-enyl-myoinositol in CH₂Cl₂ (15 ml) was added 0.2 N HCl in MeOH (15 ml, 3.00 mmol) and the reaction mixture was stirred for 1 h at 20 °C. The reaction mixture was neutralized with Et₃N and concentrated in vacuo. The residue was taken up in CH₂Cl₂, washed with H₂O, 1 M NaHCO₃ and H₂O.

The organic layer was dried over MgSO₄ and concentrated in vacuo.

Silica gel column chromatography (20 g, elution: CH₂Cl₂/MeOH, 100/0 to 95/5, v/v) of the crude product afforded pure 9 (1.31 g, 91 yield). Mp 108.5-109.5 °C (from CH₂Cl₂/hexane).

¹H-NMR (CDCl₃): delta 2.33 (d, 1H, 1-OH, exchangeable), 2.38 (d, 1H, 3-OH, exchangeable), 2.48 (d, 1H, 4-OH, exchangeable), 3.32 (dd, 1H, H-5, J_{5,6} = 9.0 Hz), 3.46 (ddd, 1H, H-3, J_{3,4} = 9.5 Hz, J_{3,OH} = 7.0 Hz),

3.57 (ddd, 1H, H-1, $J_{1,2} = 2.5$ Hz, $J_{1,OH} = 5.0$ Hz), 3.77 (dd, 1H, H-6, $J_{6,1} = 9.5$ Hz), 3.80 (s, 3H, OCH₃, pMeOBzl), 3.82 (ddd, 1H, H-4, $J_{4,5} = 9.0$ Hz, $J_{4,OH} = 2.0$ Hz), 4.01 (dd, 1H, H-2, $J_{2,3} = 2.5$ Hz), 4.71-4.95 (m, 6H, 3 x OCH₂, Bzl and pMeOBzl), 6.85-6.89 and 7.25-7.39 (m, 14H, H aromatic).

5 d) 2,6-di-O-benzyl-5-O-p-methoxybenzyl-myo-inositol 1,3,4-tris(dibenzylphosphate) (17)

A mixture of compound 9 (1.20 g, 2.50 mmol) and N,N-diisopropyl dibenzyl phosphoramidite (see K.-L. Yu and B. Fraser-Reid, *Tetrahedron Let.* 29, 979 (1988)) (3.90 g, 11.30 mmol) was coevaporated with toluene (2 x 25 ml) and dissolved in CH₂Cl₂ (30 ml). Subsequently a solution of 1H-tetrazole (1.0 g, 14.29 mmol) in CH₃CN (30 mL) was added and the reaction mixture was stirred for 15 min. ³¹P-NMR showed the presence of three peaks (delta 141.10, 141.19 and 142.55). The reaction mixture was cooled (0 °C) and tert-butyl hydroperoxide (5.65 ml) was added and stirring was continued for 45 min at 0 °C. The reaction mixture was diluted with CH₂Cl₂ and washed with H₂O, 1 M TEAB and H₂O. The organic layer was dried over MgSO₄ and concentrated in vacuo. Silica gel column chromatography (40 g, elution: hexane/EtOAc, 100/0 to 25/75, v/v) of the crude product afforded homogeneous 17 (2.74 g, 87 % yield) as an oil. ³¹P-NMR (CH₂Cl₂): delta -1.45 (1P), -1.12 (1P) and -0.82 (1P).

e) 2,6-di-O-benzyl-myo-inositol 1,3,4-tris(dibenzylphosphate) (18)

20 To a solution of compound 17 (2.52 g, 2.00 mmol) in CH₂Cl₂ (48.75 ml) was added trifluoroacetic acid (1.25 ml) and the reaction mixture was stirred for 30 min at 20 °C. The reaction mixture was diluted with CH₂Cl₂ and washed with H₂O, 1 M TEAB and H₂O. The organic layer was dried over MgSO₄ and concentrated in vacuo. Silica gel column chromatography (30 g, elution: hexane/EtOAc, 100/0 to 25/75, v/v) of the crude product afforded pure 18 (1.78 g, 78 % yield), as an oil. ³¹P-NMR (CH₂Cl₂): delta -1.30 (2P) and 0.82 (1P).

f) 2,6-di-O-benzyl-myo-inositol 5-(benzyl methylphosphonate) 1,3,4-tris(dibenzylphosphate) (20)



30 A solution of bis(1-[6-trifluoromethyl]benzotriazolyl) methyl-phosphonate in dioxane (0.2 M, 3.5 ml, 0.70 mmol) was added to compound 18 (0.40 g, 0.35 mmol), which had been dried by repeated coevaporation with pyridine. The reaction was stirred for 30 min at 20 °C. Subsequently benzyl alcohol (0.15 ml, 1.45 mmol) and N-methylimidazole (0.15 mL, 1.88 mmol) were added and the reaction mixture was stirred for another 1 h at 20 °C. After addition of 1 M TEAB the reaction mixture was diluted with CH₂Cl₂ and washed with H₂O, 1 M TEAB and H₂O. The organic layer was dried over MgSO₄ and concentrated in vacuo. Silica gel column chromatography (7.5 g, elution: hexane/EtOAc, 100/0 to 25/75, v/v) of the crude product afforded pure 20 (0.29 g, 63 % yield), as an oil. ³¹P-NMR (CH₂Cl₂): delta -1.45 (1P), -1.21(1P) and -0.70 (1P); 31.46 and 33.12 (1P, ratio, 1/3).

g) myo-inositol 5-methylphosphonate 1,3,4-trisphosphate (Na⁺ -form) (21)

40 compound 20 (215 mg, 0.16 mmol) was dissolved in a mixture of MeOH and H₂O (50 ml, 4/1, v/v) and hydrogenated over 10 % palladium on charcoal (0.20 g) at 500 kPa for 16 h at 20 °C. The solution was filtered and concentrated in vacuo (30 °C) to a small volume. After cation-exchange and lyophilization the compound 21 (102 mg, 95 % yield) was obtained, as a white solid. ³¹P-NMR (D₂O, pH = 2.00): delta 0.25 (P-1), 0.43 (P-3), 0.25 (P-4) and 31.76 (P-5). ¹H-NMR (D₂O, pH = 2.00): delta 1.48 (d, 3H, CH₃, $J_{H,P} = 1.75$ Hz), 3.90 (dd, 1H, H-6, $J_{6,1} = 10.0$ Hz), 4.08 (ddd, 1H, H-1, $J_{1,2} = 2.5$ Hz, $J_{H,P} = 8.5$ Hz), 4.18 (ddd, 1H, H-5, $J_{5,6} = 9.5$ Hz, $J_{H,P} = 9.0$ Hz), 4.20 (ddd, 1H, H-3, $J_{3,4} = 9.5$ Hz, $J_{H,P} = 9.5$ Hz), 4.44 (dd, 1H, H-2, $J_{2,3} = 2.5$ Hz), 4.50 (ddd, 1H, H-4, $J_{4,5} = 9.5$ Hz, $J_{H,P} = 9.5$ Hz).

50 Example 5

Synthesis of myo-inositol 5-ethylphosphonate 1,3,4-trisphosphate (Na⁺ -form)





To compound 18 a solution of bis(1-[6-trifluoromethyl]benzotriazolyl)ethylphosphonate was added under identical conditions as for compound 20. Following the amounts and reaction conditions as given for the compounds 20 and 21 the final product, myo-inositol 5-ethylphosphonate 1,3,4-trisphosphate (Na⁺ -form) was synthesized.

FAB-Mass-Spectrometer: Molpeak 666.

Example 6

Synthesis of myo-inositol 5-((difluoromethyl)phosphonate) 1,3,4-trisphosphate (Na⁺ -form)



a) 2,6-di-O-benzyl-myo-inositol 5-(benzyl (difluoromethyl)phosphonate) 1,3,4-tris(dibenzylphosphate) (22)

A solution of (difluoromethyl)phosphonic di(1,2,4-triazolide) in dioxane (0.2 M, 3.5 ml, 0.70 mmol) was added to compound 18 (0.40 g, 0.35 mmol), which had been dried by repeated coevaporation with pyridine. The reaction was stirred for 30 min at 20 °C. Subsequently benzyl alcohol (0.15 ml, 1.45 mmol) and N-methylimidazole (0.15 ml, 1.88 mmol) were added and the reaction mixture was stirred for another 1 h at 20 °C. After addition of 1 M TEAB the reaction mixture was diluted with CH₂Cl₂ and washed with H₂O, 1 M TEAB and H₂O. The organic layer was dried over MgSO₄ and concentrated in vacuo. Silica gel column chromatography (7.5 g, elution: hexane/ EtOAc, 100/0 to 25/75, v/v) of the crude product afforded pure 22 (0.29 g, 61 % yield), as an oil.

³¹P-NMR (CH₂Cl₂: delta -1.39, -1.15, -0.97 and 0.79 (3P); 4.35 (J_{P,F} = 95.0 Hz) and 6.86 J_{P,F} = 88.0 Hz and J_{P,F} = 102.5 Hz) (1P).

b) myo-inositol 5-((difluoromethyl)phosphonate) 1,3,4-trisphosphate (Na⁺ -form) (23)

Compound 22 (240 mg, 0.18 mmol) was dissolved in a mixture of MeOH and H₂O (50 ml, 4/1, v/v) and hydrogenated over 10 % palladium on charcoal (0.20 g) at 500 kPa for 16 h at 20 °C. The solution was filtered and concentrated in vacuo (30 °C) to a small volume. After cation-exchange and lyophilization compound 23 (114 mg, 93 % yield) was obtained, as a white solid.

³¹P-NMR (D₂O, pH = 2.00): delta 0.21 (P-1), 0.62 (P-3), 0.62 (P-4) and 4.81 (P-5). J_{P,F} = 85.5 Hz).

¹H-NMR (D₂O, pH = 2.00): delta 3.91 (dd, 1H, H-6, J_{6,1} = 10.0 Hz), 4.07 (ddd, 1H, H-1, J_{1,2} = 2.5 Hz, J_{H,P} = 8.5 Hz), 4.20 (ddd, 1H, H-5, J_{5,6} = 9.5 Hz, J_{H,P} = 9.0 Hz), 4.22 (ddd, 1H, H-3, J_{3,4} = 9.5 Hz, J_{H,P} = 9.5 Hz), 4.42 (dd, 1H, H-2, J_{2,3} = 2.5 Hz), 4.53 (ddd, 1H, H-4, J_{4,5} = 9.5 Hz, J_{H,P} = 9.5 Hz), 6.09 (ddd, 1H, CHF₂, J_{H,F} = 49.0 Hz, J_{H,P} = 24.0 Hz).

General methods and materials

Myo-inositol was purchased from Pfanstiehl Laboratories Inc. (USA).

1-Hydroxy-6-trifluoromethylbenzotriazole (see W. König and R. Geiger, Chem. Ber. 103, 788, (1970))

and 1,2,4-triazole were dried in vacuo over P_2O_5 for 70 h at 50 °C.

Triethylammonium bicarbonate buffer (TEAB, 2 M): a mixture of triethylamine (825 ml) and H_2O (2175 mL) was saturated with carbon dioxide gas at 0 °C until pH 7.0.

Column chromatography was performed on Merck Kieselgel 60 (230-400 mesh, ASTM). Melting points are uncorrected.

1H -NMR spectra were recorded on a Bruker WM-300 spectrometer, equipped with an ASPECT-2000 computer operating in the Fourier transform mode at 300 MHz.

^{31}P -NMR spectra were recorded on a Jeol JNM-FX 200 spectrometer on line with a JEC 980B computer at 50.1 and 80.7 MHz, respectively. 1H -chemical shifts are given in ppm (δ) relative to tetramethylsilane (TMS) as internal standard and ^{31}P -chemical shifts in ppm (δ) to 85 % H_3PO_4 as external standard.

Synthesis of Bis(1-[6-trifluoromethyl]benzotriazolyl) methylphosphonate


A solution of methylphosphonic dichloride (0.67 g, 5.04 mmol) in anhydrous dioxane (5 ml) was added dropwise to a stirred solution of dry 1-hydroxy-6-trifluoromethylbenzotriazole (2.05 g, 10.10 mmol) and pyridine (0.81 ml, 10.03 mmol) in anhydrous dioxane (20 mL) at 20 °C. The solution was stirred for 1 h at 20 °C and the salts were removed by filtration. The 0.2 M stock solution of bis(1-[6-trifluoromethyl]benzotriazolyl) methylphosphonate (^{31}P -NMR: δ 47.60) thus obtained could be stored for several weeks at -20 °C.

Synthesis of (Difluoromethyl)phosphonic di(1,2,4-triazolide)

A solution of (difluoromethyl)phosphonic dichloride (see D.E. Bergstrom et al., J. Org. Chem. 53, 3953 (1988)) (0.85 g, 5.03 mmol) in anhydrous dioxane (5 ml) was added dropwise to a stirred solution of dry 1,2,4-triazole (0.85 g, 12.32 mmol) and Et_3N (1.40 ml, 10.06 mmol) in anhydrous dioxane (20 ml) at 20 °C. The solution was stirred for 1 h at 20 °C and the salts were removed by filtration. The 0.2 M stock solution of (difluoro-methyl)phosphonic di(1,2,4-triazolide) (^{31}P -NMR: δ -3.01, $J_{P,F} = 106.0$ Hz) thus obtained could be stored for several weeks at -20 °C.

Claims

1. A compound of the formula (I)



in which
R1 is

- 1) phosphate
- 2) OH
- 3) 1,2 diacyl-sn-glycer-3-yl phosphate, wherein acyl is a fatty acid with 2 to 20 C-atoms and 0 to 4 C-C-double bonds, or
- 4) 1,2 dialkyl-sn-glycer-3-yl phosphate, wherein alkyl is a hydrocarbon having 2 to 20 carbon atoms and 0 to 4 C-C-double bonds

R2 is

- 1) OH, or
- 2) phosphate

R3 is

- 1) phosphate, or
- 2) OH
- 3) a radical of formula (II)



wherein

- a) Y = Z = O and X = (C₁-C₈)-alkyl
 - b) Y = Z = O and X = (C₆-C₁₈)-aryl
 - c) Y = S and Z = O and X = (C₁-C₈)-alkyl
 - d) Y = S and Z = O and X = (C₆-C₁₈)-aryl
 - e) Y = O and Z = X = (C₁-C₈)-alkyl
 - f) Y = O and Z = X = (C₆-C₁₈)-aryl
 - g) Y = S and Z = X = (C₁-C₈)-alkyl
 - h) Y = S and Z = X = (C₆-C₁₈)-aryl
 - i) Y = Z = O and X = CF₂H
 - j) Y = O and Z = X = F
 - k) Y = Z = O and X = O-(C₁-C₈)-alkyl
 - l) Y = Z = O and X = O-(C₆-C₁₈)-aryl
 - m) Y = S and Z = O and X = O-(C₁-C₈)-alkyl
 - n) Y = S and Z = O and X = O-(C₆-C₁₈)-aryl
 - o) Y = O and Z = X = O-(C₁-C₈)-alkyl
 - p) Y = O and Z = X = O-(C₆-C₁₈)-aryl
 - q) Y = S and Z = X = O-(C₁-C₈)-alkyl
 - r) Y = S and Z = X = O-(C₆-C₁₈)-aryl
- 4) a radical of formula (III)



where

- a) X = OR where R is H, (C₁-C₈)-alkyl, or (C₆-C₁₈)-aryl
- b) X = (C₁-C₈)-alkyl
- c) X = (C₆-C₁₈)-aryl
- d) X = NH-(C₁-C₈)-alkyl
- e) X = NH-(C₆-C₁₈)-aryl

- 5) a radical of formula (IV)




where

- a) X = OR where R is (C₁-C₈)-alkyl or (C₆-C₁₈)-aryl
- b) X = (C₁-C₈)-alkyl
- c) X = (C₆-C₁₈)-aryl
- d) X = NH-(C₁-C₈)-alkyl
- e) X = NH-(C₆-C₁₈)-aryl

R4 is

1) a radical of formula (V)



where

- a) Y = Z = O and X = (C₁-C₈)-alkyl
- b) Y = Z = O and X = (C₆-C₁₈)-aryl
- c) Y = S and Z = O and X = (C₁-C₈)-alkyl
- d) Y = S and Z = O and X = (C₆-C₁₈)-aryl
- e) Y = O and Z = X = (C₁-C₈)-alkyl
- f) Y = O and Z = X = (C₆-C₁₈)-aryl
- g) Y = S and Z = X = (C₁-C₈)-alkyl
- h) Y = S and Z = X = (C₆-C₁₈)-aryl
- i) Y = Z = O and X = CF₂H
- j) Y = O and Z = X = F
- k) Y = Z = O and X = O-(C₁-C₈)-alkyl
- l) Y = Z = O and X = O-(C₆-C₁₈)-aryl
- m) Y = S and Z = O and X = O-(C₁-C₈)-alkyl
- n) Y = S and Z = O and X = O-(C₆-C₁₈)-aryl
- o) Y = O and Z = X = O-(C₁-C₈)-alkyl
- p) Y = O and Z = X = O-(C₆-C₁₈)-aryl
- q) Y = S and Z = X = O-(C₁-C₈)-alkyl
- r) Y = S and Z = X = O-(C₆-C₁₈)-aryl

2) a radical of formula (VI)



wherein

- a) X = OR where R is H, (C₁-C₈)-alkyl, or (C₆-C₁₈)-aryl
- b) X = (C₁-C₈)-alkyl
- c) X = (C₆-C₁₈)-aryl
- d) X = NH-(C₁-C₈)-alkyl
- e) X = NH-(C₆-C₁₈)-aryl

3) a radical of formula (VII).



wherein

- a) X = OR where R is (C₁-C₈)-alkyl or (C₆-C₁₈)-aryl
- b) X = (C₁-C₈)-alkyl
- c) X = (C₆-C₁₈)-aryl
- d) X = NH-(C₁-C₈)-alkyl
- e) X = NH-(C₆-C₁₈)-aryl

and the physiologically tolerated salts thereof.

2. A compound of the formula (I) as claimed in claim 1, in which
- R1 is phosphate or 1,2-diacyl-sn-glycer-3-yl-phosphate, wherein acyl is a fatty acid radical having 2 to 20 carbon atoms and 0 to 4 C-C-double bonds,
 - R2 is OH or phosphate,
 - R3 is OH or phosphate,
 - R4 is a radical of formula (V)



wherein $Y = Z = O$ and $X = (C_1-C_8)$ -alkyl,
and the physiologically tolerated salts thereof.

3. A compound of formula (I) as claimed in claim 2, in which
- R4 is a radical of the formula (V) wherein X is methyl, ethyl or propyl.
4. Process for the preparation of a compound of the formula (I) as claimed in one of the claims 1 to 3 starting with a myo-inositol derivative.
5. A compound of the formula (I) as claimed in one of the claims 1 to 3 for the use as medicine.
6. A compound of the formula (I) as claimed in one of the claims 1 to 3 for the use as a medicine for the treatment of cardiovascular diseases.
7. A pharmaceutical composition comprising an effective amount of a compound as claimed in claim 1, or a physiologically tolerated salt thereof, and a physiologically acceptable vehicle.



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number

EP 92 11 0563

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. CL.5)
Y	EP-A-0 262 227 (MITSUI TOATSU CHEMICALS) * the whole document * ---	1-7	C07F9/40 A61K31/66 C07F9/117
Y	US-A-4 515 722 (SHU S. YANG) * the whole document * ---	1-7	
Y	EP-A-0 269 105 (SIREN, MATTI) * the whole document * ---	1-7	
Y	TETRAHEDRON, (INCL. TETRAHEDRON REPORTS) vol. 47, no. 26, 24 June 1991, OXFORD GB pages 4709 - 4722; C.E. DREEF ET AL.: 'synthesis of racemic 3-methylphosphonate analogues of myo-inositol 3,4-bis- and 1,3,4-trisphosphate' * the whole document * -----	1-7	
			TECHNICAL FIELDS SEARCHED (Int. CL.5)
			C07F
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 22 SEPTEMBER 1992	Examiner BESLIER L.M.
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure F : intermediate document		I : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons ----- & : member of the same patent family, corresponding document	

THIS PAGE BLANK (USPTO)